

Ethnobotany. Lecture 16

Alexey Shipunov

Minot State University

February 23th, 2011

Outline

- 1 Fruits: Introduction
- 2 Fruit plants
 - Rosaceae fruits



Outline

- 1 Fruits: Introduction
- 2 Fruit plants
 - Rosaceae fruits

Organic acids

- Malic (*Pyrus malus*, apple and other Rosaceae fruits)
- Citric (*Citrus* fruits etc.)
- Tartaric (e.g., in wine)
- Are good antioxidants

Citric and malic acids




Fruit vitamins

- Vitamin C (ascorbic acid)
- Pro-vitamin A (β -carotene)
- Other carotenes (lycopene etc.)

Specific components which are restricted to few species

- Lipids
- Starch
- Gums, mucilages, pectins
- Astringent chemicals (e.g., tannic acid)
- Aroma compounds
- Other secondary* metabolites (latex, alkaloids, glycosides)

Tannic acid



Rosaceae in general

- Medium-sized family ($\approx 3,000$ species) of small trees, shrubs and herbs from subtropical and temperate regions
- Flower contains numerous stamens (secondary multiplied), free pistils and hypanthium
- Fruit is mostly fleshy

Rosaceae groups

- Rosoideae—herbs or shrubs, leaves often compound, receptacle large, fruit aggregate
- Spiraeoideae—shrubs or trees, leaves simple, receptacle small, fruit often monomerous
- Maloideae—trees, leaves simple, receptacle and pistils fused

Rosaceae with multiple (aggregate) fruits

- Most primitive group
- *Rosa* is ornamental and medicine plant
- *Rubus* and *Fragaria* are also widely cultivated

Rubus

- Biennial semi-shrubs, sometimes herbs
- Multiple wild species, only two are widely cultivated: raspberry (*Rubus idaea*) and blackberry, *Rubus caesius* forms and hybrids

Rubus features

- Two aboveground stem types: primocane and floricanes, plus underground rhizomes
- Tangled genetic systems: apomixis, polyploidy and even permanent pentaploidy ($2n = 35$)
- Fruits contain (among other) salicylic acid and different antioxidants

Blackberries on the different stages or ripening



Fragaria × *ananassa*, strawberry

- Octoploid ($2n = 56$) hybrid species of two other octoploid strawberries, *Fragaria virginiana* from North America and *F. chiloensis* from Chile.
- Garden hybrid, first occurrences are from ≈ 1740
- Herb with runner stems and accessory multiple nut fruit (the edible part is a receptacle)

Strawberry features

- Susceptible to multiple diseases, often cultivated in semi-artificial conditions as plasticulture
- Cultivated as annual or perennial
- Long-day cultivars flower early in May and capable to produce fruits in June

Plasticulture of strawberries



Summary

- Rosaceae is one of the most important temperate fruit families
- Most of Rosaceae cultivated fruits are result of long selection involved multiple hybridization

For Further Reading



A. Shipunov.

Ethnobotany [Electronic resource]. 2011—onwards.

Mode of access:

http://ashipunov.info/shipunov/school/biol_310



P. M. Zhukovskij.

Cultivated plants and their wild relatives [Electronic resource].

Commonwealth Agricultural Bureaux, 1962.

Mode of access:

[http://ashipunov.info/shipunov/school/biol_310/
zhukovskij1962_cultivated_plants.pdf](http://ashipunov.info/shipunov/school/biol_310/zhukovskij1962_cultivated_plants.pdf)

Pages 28–74.