

Biometry. Lecture 15

Alexey Shipunov

Minot State University

March 30, 2016



1 Two-dimensional statistics

2 Effect sizes

- Test for tables: chi-squared



- 1 Two-dimensional statistics
- 2 Effect sizes
 - Test for tables: chi-squared



```
> setwd("<working folder>")  
or  
"Change dir"  
in menu!
```

On Mac, be sure that startup option is working: `getwd()`
(`getwd()` checks if R is in working folder, `dir()` checks the folder content)



Quiz question (10 points!)



Quiz question (10 points!)

These are points from the first and second exam in one small class:

63, 72, 77, 76, 67, 56, 55, 51, 77, 64

and

87, 86, 76, 79, 54, 60, 97, 80, 73, 97

Both exams were equivalent. Provide a statistical support for the hypothesis that second exam went better. Report commands and all values which support your conclusion.



Effect sizes: parametric and non-parametric

- p-values do not show the extent of differences
- Means and alike are not working because data could intersect
- Consequently, effect sizes were invented. They allow to understand how big is a difference
- R library `effsize` provides parametric Cohen D and non-parametric Cliff's Delta effect sizes



Effect sizes: examples

```
> library(effsize)
> a <- 1:9
> b <- rep(5, 9)
> cliff.delta(a, b) # no effect
> cohen.d(extra ~ group, data=sleep)
```

In the `sleep` example, effect size is present even if p-value is not good enough!



Effect sizes

Test for tables: chi-squared



Contingency tables

- Secondary data: counts
- May be created from any categorical variable, or from measurement variable after cutting



table() function

```
> with(airquality, table(cut(Temp, quantile(Temp)), Month))  
> d <- factor(rep(c("A", "B", "C"), 10))  
> is.na(d) <- 3:4  
> table(d, exclude=NULL)
```



Graphical representation of tables

```
> Titanic # this is a multidimensional table  
> ftable(Titanic, row.vars = 1:3)  
> margin.table(Titanic, c(1, 4)) # make 2 dimensions  
> mosaicplot(margin.table(Titanic, c(1, 4)))
```



- Chi-squared test checks the null if *variables in the table are distributed independently* (non-accordingly) between cells.
- Alternative hypothesis is that association between variables exists.



Chi-squared test

```
> HairEyeColor # multidimensional table  
> margin.table(HairEyeColor, c(1, 2)) # hairs and eyes  
> chisq.test(margin.table(HairEyeColor, c(1, 2)))  
> margin.table(HairEyeColor, c(2, 3)) # eyes and sex  
> chisq.test(margin.table(HairEyeColor, c(2, 3)))
```



Association plot

```
> assocplot(margin.table(HairEyeColor, c(1, 2)))
```

Association plots show positive and negative association between factors in the table. The key thing is the **asymmetry between the squares**.



Finishing...

Save your commands!

`(savehistory(<today's date>.r)` or File -> Save as... on
Mac)



Summary: most important commands

- `response ~ factor`—if factor has exactly two levels, this is a model formula for two-sample test
- `table()`—creates contingency tables
- `chisq.test()`—test for independence of rows and columns



For Further Reading



A. Shipunov.

Biometry [Electronic resource].

2012—onwards.

Mode of access:

http://ashipunov.info/shipunov/school/biol_240



A. Shipunov, and many others.

Visual statistics. Use R!

2016—onwards.

Mode of access: http://ashipunov.info/shipunov/school/biol_240/en/visual_statistics.pdf

